



## Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin

ICAR Central Coastal Agricultural Research Institute, Ela  
North Goa



### Agromet Advisory Bulletin

Date : 08-02-2022

Weather Forecast of NORTH-GOA(Goa) Issued On : 2022-02-08(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-02-09	2022-02-10	2022-02-11	2022-02-12	2022-02-13
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	31.0	31.0	32.0	33.0	32.0
Tmin(°C)	20.0	20.0	20.0	21.0	22.0
RH-I(%)	88	70	63	51	52
RH-II(%)	45	34	29	25	22
Wind Speed(kmph)	9.2	9.5	8.3	8.0	7.6
Wind Direction(Degree)	288	292	90	79	90
Cloud cover(octa)	1	2	2	1	0

#### Weather Summary/Alert:

• Weather is very likely to be dry for next 5 days • No large change in minimum temperature for 2 days and gradual rise by 1-2°C in the subsequent 3 days • Maximum & minimum temperatures are likely to be around 30°C & 20°C respectively Week 1 (04th Feb to 10th Feb) • Rainfall amount is very likely to be normal over the district • Maximum temperature is likely to be normal • Minimum temperature is likely to be normal Week 2 (11th Feb to 17th Feb) • Rainfall amount is very likely to be normal over the district • Maximum temperature is likely to be normal • Minimum temperature is likely to be normal

#### General Advisory:

- Keep the orchards clean, by weeding and removing debris
- Mulching of tree basins should be done to conserve soil moisture

### SMS Advisory:

- Spraying of insecticides in mango should be done before 9am or after 4pm in order to save non-target pollinators

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
RICE	<ul style="list-style-type: none"> <li>• Possibility of leaf folder in rabi paddy, hence early clipping of infested leaf tips along with removal of other weeds is recommended</li> <li>• Foliar sprays with Chlorpyrifos 2.5 ml/litre is recommended</li> <li>• Application of second dose of fertilizer (Urea – 54kg/ha) can be done at tillering stage</li> <li>• Maintain the water level in the paddy fields</li> </ul>
GROUNDNUT	<ul style="list-style-type: none"> <li>• Earthing up is to be done within 40-45 days after sowing as it helps for the penetration of pegs in the soil and also facilitates for increased pod development</li> <li>• Provide irrigation at 10 to 15 days interval</li> <li>• To control the aphids, thrips and leaf miners spraying of spinosad @ 0.2 ml/litre of water can be done or apply NSKE 5% (neem seed kernel extract)</li> <li>• Yellow sticky traps can be kept in the field for better monitoring of these sucking pests</li> </ul>
CASHEW	<ul style="list-style-type: none"> <li>• Due to the dry weather forecast for the next 5 days, irrigation can be given at 15 days interval</li> <li>• Farmers are advised to look upon the cashew orchards for the Tea mosquito bug incidence and apply neem-based insecticides</li> </ul>

### Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
COWPEA/ LOBIA	<ul style="list-style-type: none"> <li>• Foliar fertilization may be practiced with 2% DAP (200g dissolved in 10 litres of water) at flowering (40- 45 DAS) which considerably increases 20 to 35 % of yield</li> <li>• Farmers are suggested to take up weeding and earthing up practices</li> <li>• Provide irrigation at 7 to 10 days interval</li> <li>• To control the aphids and thrips spraying of spinosad @ 0.2 ml/litre of water can be done or apply NSKE 5% (neem seed kernel extract)</li> <li>• Yellow sticky traps can be kept in the field for better monitoring of these sucking pests</li> </ul>

<b>Horticulture(Varieties)</b>	<b>Horticulture Specific Advisory</b>
CHILLI	<ul style="list-style-type: none"> <li>• Provide irrigation at regular intervals</li> <li>• Take up intercultural operations and keep the fields weed-free</li> <li>• Take up control measures against chilli leaf curl diseases</li> <li>• Spraying of Imidacloprid @ 0.3ml per litre of water can be done</li> <li>• Yellow sticky traps can be placed in the field for better monitoring of sucking pests</li> </ul>
MANGO	<ul style="list-style-type: none"> <li>• Spraying of 0.3 ml Imidacloprid per litre of water is suggested to control hoppers</li> <li>• Spraying should be done before 9am or after 4pm in order to save non-target pollinators</li> <li>• Provide irrigation for below 5 years old trees at 10-15 days interval</li> </ul>
ARECANUT	<ul style="list-style-type: none"> <li>• Due to dry weather forecast for next 5 days , irrigation can be given to arecanut palms in sandy soils at 4 days interval in basin method or daily through drip irrigation</li> <li>• Keep the orchards clean, by weeding and removing debris</li> <li>• Mulching of tree basins should be done to conserve soil moisture</li> <li>• Protect the palms from sun-scorch</li> <li>• Harvesting and drying of ripe nuts</li> </ul>
COCONUT	<ul style="list-style-type: none"> <li>• Due to dry weather forecast for next 5 days , irrigation can be given to coconut palms in sandy soils at 4 days interval in basin method or daily through drip irrigation</li> <li>• To control mites, spraying of 2% neem-garlic emulsion or azadirachtin 10000 ppm @0.004% is recommended</li> <li>• To control white flies , application of 1% starch solution on leaflets to flake out the sooty moulds</li> <li>• In severe case, spray neem oil 0.5% and no insecticide is recommended</li> <li>• Installation of yellow sticky traps on the palm trunk to trap adult whiteflies</li> </ul>

### Live Stock Specific Advisory:

<b>Live Stock(Varieties)</b>	<b>Live Stock Specific Advisory</b>
PIG	<ul style="list-style-type: none"> <li>• There is a forewarning of very high risk of Peste-des-Petitis Ruminants (PPR) and Swine Fever (SF) in livestock for North Goa district in February 2022</li> <li>• Cleaning and disinfection of pig sheds and goat farms should be done</li> <li>• Vaccination should be followed strictly</li> </ul>