



# Gramin Krishi Mausam Sewa

District Level Agromet Advisory Bulletin  
ICAR Central Coastal Agricultural Research Institute, Ela  
North Goa



## Agromet Advisory Bulletin

Date : 20-09-2022

Weather Forecast of NORTH-GOA(Goa) Issued On : 2022-09-20(Valid Till 08:30 IST of the next 5 days)

Parameter	2022-09-21	2022-09-22	2022-09-23	2022-09-24	2022-09-25
Rainfall(mm)	2.0	2.0	4.0	5.0	5.0
Tmax(°C)	31.0	31.0	31.0	32.0	32.0
Tmin(°C)	23.0	23.0	23.0	23.0	23.0
RH-I(%)	91	92	92	92	92
RH-II(%)	64	64	64	63	67
Wind Speed(kmph)	7.7	6.8	6.6	6.6	9.1
Wind Direction(Degree)	289	292	112	248	296
Cloud cover(octa)	6	8	7	1	2

### Weather Summary/Alert:

• Light to moderate rain very likely at many places on 20 & 21 september 2022 • Light rain very likely at a few places on 22nd , 23rd & 24th september 2022 • No weather warnings for next 5 days • Maximum & minimum temperatures are likely to be around 31°C & 23°C respectively Week 1 ( Sep 16th – Sep 22nd ) • Rainfall is very likely to be normal • Maximum temperature is likely to be 0-2°C lower than normal • Minimum temperature is likely to be 0-2°C lower than normal Week 2 ( Sep 23rd – Sep 29th ) • Rainfall is very likely to be normal • Maximum temperature is likely to be 0-2°C lower than normal • Minimum temperature is likely to be 0-2°C higher than normal

### General Advisory:

• Make arrangements to drain out excess water • Take control measures pest and diseases during clear weather (no rainfall at least 5-6 hrs)

### SMS Advisory:

• Application of fertilizers can be done for the 5 years above coconut and arecanut palms during last week of September

### Crop Specific Advisory:

Crop(Varieties)	Crop Specific Advisory
-----------------	------------------------

Crop(Varieties)	Crop Specific Advisory
RICE	<ul style="list-style-type: none"> <li>• Farmers are suggested to maintain the water level in the field</li> <li>• Possibility of incidence of leaf folder and stem borer, hence keep the field and bunds clean by removing weeds. Avoid use of excessive fertilizers. Pull out and destroy the affected tillers (dead hearts). Need based spraying of chlorpyrifos @ 2 ml/litre during clear weather (no rainfall at least 5-6 hrs)</li> <li>• Possibility of incidence of gundi bug. Hence Egg masses, nymph and adults can be collected (hand or using hand nets) and destroyed.</li> <li>Removal of grasses near the paddy fields. Dusting Malathion 5 D @ 25 kg/ha starting from field edges to the centre in a circular manner during clear weather (no rainfall at least 5-6 hrs)</li> </ul>
CASHEW	<ul style="list-style-type: none"> <li>• Due to humidity and rainfall, possibility of incidence of cashew stem and root borer</li> <li>• Hence control measure can be taken up during clear weather (no rainfall at least 5-6 hrs)</li> <li>• Treat the infested trees of the trunk region up to one meter height from the ground level and on exposed roots with Fipronil @ 2 ml / litre of water</li> </ul>

#### Horticulture Specific Advisory:

Horticulture(Varieties)	Horticulture Specific Advisory
MANGO	<ul style="list-style-type: none"> <li>• Possibility of anthracnose in mango trees</li> <li>• Hence control measure can be taken up during clear weather (no rainfall at least 5-6 hrs)</li> <li>• Spraying of carbendazim @ 2 g / liter of water is recommended to protect the new flushing in mango from the incidence of anthracnose</li> </ul>
COCONUT	<ul style="list-style-type: none"> <li>• Circular basins of 1.8m in radius and 25 cm depth may be dug and green leaf or compost or farm yard manure may be spread in the basin</li> <li>• Apply DAP-860 g + Urea-270g + MOP-1200g + Zinc sulfate heptahydrate-95g + Borax-70g per tree during last week of September for 5 years above palm</li> <li>• Possibility of incidence of but rot. Hence control measure can be taken up during clear weather (no rainfall at least 5-6 hrs)</li> <li>• Spraying of 1% Bordeaux mixture is recommended</li> <li>• Possibility of incidence of rugose spiralling whitefly.</li> <li>• In juvenile palms, spraying of water with jet speed could dislodge the whitefly and reduce the feeding as well as breeding potential of the pest</li> <li>• Installation of yellow sticky traps</li> <li>• In severe cases, spray only neem oil 0.5% or NSKE 5% and avoid spraying any form of insecticides.</li> </ul>
ARECANUT	<ul style="list-style-type: none"> <li>• Apply DAP-385 g + Urea-70g + MOP-200g + Zinc sulfate heptahydrate-95g + Borax-70g per tree during last week of September for 5 years above tree</li> <li>• Possibility of incidence of fruit rot . Hence control measure can be taken up during clear weather (no rainfall at least 5-6 hrs)</li> <li>• Spraying of 1% Bordeaux mixture is recommended</li> </ul>

#### Live Stock Specific Advisory:

Live Stock(Varieties)	Live Stock Specific Advisory
COW	<ul style="list-style-type: none"> <li>• Supplementation of Perrineal Napier grass, Subabul, and Moringa leaves with traditional feed for enriching the quality of ration</li> <li>• Deworming of all the adult stock with broad spectrum anthelmintics like Albendazole (Dose: 10 mg/ kg body weight) during last week</li> <li>• FMD vaccine during month of September</li> </ul>

#### Poultry Specific Advisory:

Poultry(Varieties)	Poultry Specific Advisory
--------------------	---------------------------

Poultry(Varieties)	Poultry Specific Advisory
BIRD	<ul style="list-style-type: none"> <li>• Inclusion of different cereal sources such as maize/wheat/barley/ragi/broken rice at 50%,</li> <li>• Protein sources like soyabean meal/groundnut meal/sunflower meal/linseed meal upto 20-25%,</li> <li>• Rice polish/rice bran/wheat bran upto 10-20%</li> <li>• Gram chuni at 5-10%, coconut meal and leafy meals 5% each or molasses 5% and vitamin and mineral mix 2% totalling to 100 in different combinations can be used for making poultry feed</li> <li>• Deworming of backyard poultry in deep litter housing should be done at 8 weeks and 16/18 weeks then once in a month with suitable anthelmintics</li> </ul>